Journal of Immuological Methods, 122 (1989) 298–299
Elsevier

Contents to volume 122 (1989)

Number 1 – 15 August 1989

Immunological standardization

Designation of antibodies and their derivatives. Suggestions for a general nomenclature – A discussion document
H. Baumgarten and K. Kürzinger (F.R.G.) 1

Research reports

MicroELISA method for the determination of thymosin β_9 discriminating between thymosin β_9 and the structurally closely related thymosin β_4
M. Mihelič, H. Kalbacher, E. Hannappel and W. Voelter (F.R.G.) 7

Comparison of fluorochrome-labeled and ^{51}Cr-labeled targets for natural killer cytotoxicity assay
W.G. Wierda, D.S. Mehr and Y.B. Kim (U.S.A.) 15

A new tool for the serodiagnosis of acute *Plasmodium falciparum* malaria in individuals with primary infection
K. Früh, H.-M. Müller, H. Bujard and A. Crisanti (F.R.G.) 25

Biotinylation of human interleukin-2 for flow cytometry analysis of interleukin-2 receptors
S. Taki, T. Shimamura, M. Abe, T. Shirai and Y. Takahara (Japan) 33

Analysis of anti-insulin antibodies using time-resolved fluoroimmunoassay
Q. Vos and R. Benner (The Netherlands) 43

Determination of the components of immune complexes made in vitro with antigens derived from *Pseudomonas aeruginosa*
G. Kronborg, A. Fomsgaard, G.H. Shand and N. Høiby (Denmark) 51

Sensitive time-resolved fluoroimmunoassay of nerve growth factor and the disappearance of nerve growth factor from rat pheochromocytoma PC12 cell culture medium
U. Arumäe, T. Neuman, R. Sinijärv and M. Saarma (U.S.S.R.) 59

A novel and simple procedure for determining T cell epitopes in protein antigens
S. Demotz, P. Matricardi, A. Lanzavecchia and G. Corradin (U.S.A., Switzerland) 67

Rapid isolation and characterization of native mouse complement components C3 and C5
C.W. Van den Berg, H. Van Dijk and P.J.A. Capel (The Netherlands) 73

Measurement of binding specificity between T cell lymphomas and murine leukemia viruses
H.C. O'Neill and I.L. Weissman (U.S.A.) 79

A dual microtiter plate (192 sample) luminometer employing computer-aided single-photon imaging applicable to cellular luminescence and luminescence immunoassay
F.-E. Maly, M. Vittoz, A. Urwyler, K. Koshikawa, L. Schleinkofer and A.L. De Weck (Switzerland, F.R.G.) 91

Purification of human blood eosinophils by negative selection using immunomagnetic beads
T.T. Hansel, J.D. Pound, D. Pilling, G.D. Kitas, M. Salmon, T.A. Gentle, S.S. Lee and R.A. Thompson (U.K.) 97

Separation of active and inactive forms of the third component of human complement, C3, by fast protein liquid chromatography (FPLC)
P. Sánchez-Corral, L.C. Antón, J.M. Alcolea, G. Marqués, A. Sánchez and F. Vivanco (Spain) 105

The FITC-anti-FITC system is a sensitive alternative to biotin-streptavidin in ELISA
I.J. Harmer and D. Samuel (U.K.) 115

An enzyme-linked immunosorbent assay (ELISA) for the measurement of serum antibodies to growth hormone. The Wessex Growth Study
T. Wilkin, C. Casey, A. Tuck and P. Englefield (U.K.) 123

Loss of antigens from immunoblotting membranes
N. DenHollander and D. Befus (Canada) 129

Production of large numbers of hybridomas producing monoclonal antibodies against rat IgE using mast cell-deficient w/w^v and sl/sl^d strains of mice
B.J. Rup (U.S.A.) 137

A miniaturised method for immunoglobulin allotype serology
N.T. Young, J. Street and C. Darke (U.K.) 143

Technical note

The use of foetal calf serum as a blocking agent in double immunoenzyme-cytochemistry labelling with mouse monoclonal antibodies
P.A. Lalor and P.A. Revell (U.K.) 149

0022-1759/89/$03.50 © 1989 Elsevier Science Publishers B.V. (Biomedical Division)

Journal of Immuological Methods, 122 (1989) 298–299
Elsevier

Contents to volume 122 (1989)

Number 1 – 15 August 1989

Immunological standardization

Designation of antibodies and their derivatives. Suggestions for a general nomenclature – A discussion document
H. Baumgarten and K. Kürzinger (F.R.G.) 1

Research reports

MicroELISA method for the determination of thymosin β_9 discriminating between thymosin β_9 and the structurally closely related thymosin β_4
M. Mihelič, H. Kalbacher, E. Hannappel and W. Voelter (F.R.G.) 7

Comparison of fluorochrome-labeled and ^{51}Cr-labeled targets for natural killer cytotoxicity assay
W.G. Wierda, D.S. Mehr and Y.B. Kim (U.S.A.) 15

A new tool for the serodiagnosis of acute *Plasmodium falciparum* malaria in individuals with primary infection
K. Früh, H.-M. Müller, H. Bujard and A. Crisanti (F.R.G.) 25

Biotinylation of human interleukin-2 for flow cytometry analysis of interleukin-2 receptors
S. Taki, T. Shimamura, M. Abe, T. Shirai and Y. Takahara (Japan) 33

Analysis of anti-insulin antibodies using time-resolved fluoroimmunoassay
Q. Vos and R. Benner (The Netherlands) 43

Determination of the components of immune complexes made in vitro with antigens derived from *Pseudomonas aeruginosa*
G. Kronborg, A. Fomsgaard, G.H. Shand and N. Høiby (Denmark) 51

Sensitive time-resolved fluoroimmunoassay of nerve growth factor and the disappearance of nerve growth factor from rat pheochromocytoma PC12 cell culture medium
U. Arumäe, T. Neuman, R. Sinijärv and M. Saarma (U.S.S.R.) 59

A novel and simple procedure for determining T cell epitopes in protein antigens
S. Demotz, P. Matricardi, A. Lanzavecchia and G. Corradin (U.S.A., Switzerland) 67

Rapid isolation and characterization of native mouse complement components C3 and C5
C.W. Van den Berg, H. Van Dijk and P.J.A. Capel (The Netherlands) 73

Measurement of binding specificity between T cell lymphomas and murine leukemia viruses
H.C. O'Neill and I.L. Weissman (U.S.A.) 79

A dual microtiter plate (192 sample) luminometer employing computer-aided single-photon imaging applicable to cellular luminescence and luminescence immunoassay
F.-E. Maly, M. Vittoz, A. Urwyler, K. Koshikawa, L. Schleinkofer and A.L. De Weck (Switzerland, F.R.G.) 91

Purification of human blood eosinophils by negative selection using immunomagnetic beads
T.T. Hansel, J.D. Pound, D. Pilling, G.D. Kitas, M. Salmon, T.A. Gentle, S.S. Lee and R.A. Thompson (U.K.) 97

Separation of active and inactive forms of the third component of human complement, C3, by fast protein liquid chromatography (FPLC)
P. Sánchez-Corral, L.C. Antón, J.M. Alcolea, G. Marqués, A. Sánchez and F. Vivanco (Spain) 105

The FITC-anti-FITC system is a sensitive alternative to biotin-streptavidin in ELISA
I.J. Harmer and D. Samuel (U.K.) 115

An enzyme-linked immunosorbent assay (ELISA) for the measurement of serum antibodies to growth hormone. The Wessex Growth Study
T. Wilkin, C. Casey, A. Tuck and P. Englefield (U.K.) 123

Loss of antigens from immunoblotting membranes
N. DenHollander and D. Befus (Canada) 129

Production of large numbers of hybridomas producing monoclonal antibodies against rat IgE using mast cell-deficient w/w^v and sl/sl^d strains of mice
B.J. Rup (U.S.A.) 137

A miniaturised method for immunoglobulin allotype serology
N.T. Young, J. Street and C. Darke (U.K.) 143

Technical note

The use of foetal calf serum as a blocking agent in double immunoenzyme-cytochemistry labelling with mouse monoclonal antibodies
P.A. Lalor and P.A. Revell (U.K.) 149

0022-1759/89/$03.50 © 1989 Elsevier Science Publishers B.V. (Biomedical Division)

299

Announcement 151

Notes to authors 152

Number 2 – 1 September 1989

Immunological standardization
Cytokine standardization 153

Research reports
Specific immunocytochemical visualization of phosphatidylserine
L. Maneta-Peyret, G. Freyburger, J.-J. Bessoule and C. Cassagne (France) 155
Whole-blood lymphocyte cultures
E. Bloemena, M.T.L. Roos, J.L.A.M. Van Heijst, J.M.J.J. Vossen and P.T.A. Schellekens (The Netherlands) 161
Latex agglutination assay for human anti-*Brucella* IgM antibodies
C.L. Cambiaso and J.N. Limet (Belgium) 169
Quantitation of effector-target affinity in the human NK cell and K562 tumor cell system
P. Garcia-Peñarrubia and A.D. Bankhurst (U.S.A.) 177
2- and 4-iodinated estriol as indicator ligands for estriol radioimmunoassays with anti-estriol-C6 conjugate antiserum
K. Gloning, L. Dibbelt and E. Kuss (F.R.G.) 185
The transport and culture conditions for optimum transformation responses of wildfowl lymphocytes to mycobacterial antigens
R.L. Cromie, J.L. Stanford, M.J. Brown and D.J. Price (U.K.) 195
High yields of stable transformants by hypo-osmolar plasmid electroinjection
R. Däumler and U. Zimmermann (F.R.G.) 203
Demonstration of IL-1α and IL-1β secretion by the monocytic leukemia cell line, THP-1
E.V. Gaffney, C.R. Stoner, S.E. Lingenfelter and L.A. Wagner (U.S.A.) 211
Specific detection of *Haemophilus influenzae* type b lipooligosaccharide by a polymyxin B monoclonal antibody assay
J. Mertsola, O. Ramilo, X. Sáez-Llorens, M.S. Hanson, G.H. McCracken, Jr. and E.J. Hansen (U.S.A.) 219
Improved anti-tumor reactivity with monoclonal anti-idiotypic antibody conjugated to syngeneic mouse red blood cells
W.-Z. Wei, S. Fisher and H. Kohler (U.S.A.) 227
The kinetics of interaction between lymphocytes and magnetic polymer particles
D. Pilling, G.D. Kitas, M. Salmon and P.A. Bacon (U.K.) 235
A rapid mechanical vibration procedure for purifying mature (Ia positive) columnar epithelial cells from the mouse small intestine
D. Kaiserlian, C. Roche, M. Blanc and J.P. Revillard (France) 243
Comparison of three different activation methods for coupling antibodies to magnetisable cellulose particles
I.H. Al-Abdulla, G.W. Mellor, M.S. Childerstone, A.M. Sidki and D.S. Smith (U.K.) 253
A simplified solid-phase assay for the quantitation of native membrane proteins. Application to the measurement of EGF receptor induction by dexamethasone
M. Von Knebel Doeberitz, H. Drzonek, S. Koch and C.-M. Becker (F.R.G.) 259
A simple immunofiltration assay for mucins secreted by a human colonic epithelial cell line
I. Roumagnac and C.L. Laboisse (France) 265
Preparation of nanoparticles conjugated with enzyme and antibody and their use in heterogeneous enzyme immunoassays
K.G.I. Nilsson (Sweden) 273
Measurement of plasma glycoalbumin levels with a monoclonal antibody based ELISA
M.P. Cohen and E. Hud (U.S.A.) 279
Preparation and properties of gel-peptide immunoadsorbents
A. Chersi, E. Ruocco and E. Muratti (Italy) 285
Triton X-100 eliminates plasma proteins interference in a radioimmunoassay for luteinizing hormone-releasing hormone (LHRH) and LHRH analogues
E. Ezan, K. Drieu and F. Dray (France) 291

Announcement 297

Contents to volume 122 (1989) 298

Notes to authors 300